

CITY AND COUNTY OF SAN FRANCISCO PUBLIC HEALTH LABORATORY

101 Grove Street, Room 419 San Francisco, CA 94102 Tel: (415) 554-2800 Fax: (415) 431-0651 Godfred Masinde, PhD., Lab Director CLIA ID # 05D0643643

Test Order

AFB (Acid Fast) Culture

Symonym (a)	Mycobacteria culture, detection and recovery of Mycobacterium sp.
Synonym(s) Methodology	Mycobacteria culture, detection and recovery of <i>Mycobacterium sp.</i> Mycobacteria culture using liquid (BACTEC MGIT 960) and solid growth media (Middlebrook agar)
Acceptable Specimen Type(s) for Testing	Sputum (expectorated or induced); urine (first morning collection)
Transport / Collection Medium	Sputum: sterile collection cup; urine: large sterile plastic container
Storage and Preservation of Specimen	Refrigerate at 4°C after collection and deliver to the laboratory within 3 days.
Minimum Volume Required	3 – 10 mL (1 mL is acceptable) sputum; >50 mL urine
Additional Collection Instructions	It is ideal to have at least three consecutive sputum specimens obtained, each collected within 24 hours and at least one being an early morning specimen. Have the patient rinse mouth with water before collecting the specimen to minimize contamination with food particles, mouth wash, or oral drugs which might inhibit the growth of mycobacteria.
Additional Required Information	Label specimen with patient name, date of birth, medical record number and date of collection; Label requisition with patient name, date of birth, address, medical record number, gender, date of collection, submitter name, requesting clinician, specimen source and test request.
Send Out?	No
Turnaround Time	6 weeks
Testing Restrictions	Rejection Criteria: Leaky specimens; No label on the specimen; Requisition and specimen labels do not match; Quantity not sufficient for testing; Specimen is too old for testing; Specimen was not stored properly
Requisition Form(s)	http://sfcdcp.org/document.html?id=1035
Limitations / Notes / Disclaimers	Recovery of mycobacteria species is dependent upon quality of specimen and collection. It is critical to follow the collection procedure to ensure inhibitory substances are removed from the mouth.

Updated 5/04/2017